

stemtrac®

What is stemtrac®?

Tumors release cells into the surrounding tissue and into the blood. They are called circulating epithelial tumor cells (CETCs/CTCs). Amongst them are the so-called **circulating cancer stem cells**, which can be identified with **stemtrac®**. During a period of up to **21 days**, the cancer stem cells grow into **tumorspheres** in vitro. A tumorsphere is a **spherical structure** that results from the cell division of a cancer stem cell¹.

Clinical Relevance¹

The more stemtrac® Tumorspheres found, the more aggressive the tumor and the higher the risk of metastasis.

- Metastasized patients have more tumorspheres than non-metastasized patients.
- The number of tumorspheres can be used as a biomarker for the presence of already existing metastases.
- No tumorsphere growth was observed in subjects without diagnosed cancer.

Application of stemtrac®

The number of **stemtrac® Tumorspheres** can be used in combination with **maintrac®** to monitor the activity of the remaining tumor burden:


- After the end of therapy to estimate the aggressiveness and metastatic risk of the remaining tumor cells.
- In complete remission when CETCs are increasing.
- In disease progression when CETCs are low or undetectable (loss of EpCAM expression, i.e. dedifferentiation of tumor cells).

Please note that in case of chemotherapy, blood sampling for stemtrac® tumorspheres should be performed 2-3 weeks after administration of the cytostatic drugs.

Innovative Laboratory Diagnostics of Circulating Cancer Stem Cells Before, During and After Cancer Therapy

stemtrac® Quality Features

- **Detection** of circulating cancer stem cells without enrichment steps^{1,2}
- **Quantitative** determination of circulating cancer stem cells from peripheral blood^{1,2}
- Highly sensitive and reproducible^{1,2}
- Performed in a DIN EN ISO 15189 certified laboratory, accredited by DAkkS (ILAC approved)³



The pictures show the development of tumorspheres of circulating cancer stem cells from peripheral blood.

Additional examinations

- maintrac® Cell Counting
- maintrac® Therapeutic Substance Testing
- maintrac® Therapy Relevant Tumor Cell Characteristics

Requisition

Shipping boxes including the lab request form can be ordered free of charge online at:

www.maintrac.de/en/order/order-maintrac-boxes


Only 15 ml EDTA blood is required for the examination.

Transmission of Results

The results are usually sent **digitally** (DSGVO-compliant) or **by post** within three weeks.

0241106-FTS-EN-WM

Your competent partner in



Laboratory Dr. Pachmann Kurpromenade 2 95448 Bayreuth Phone: +49 921 850 200 E-mail: mail@laborpachmann.de www.laborpachmann.de

www maintrac de

Costs

At present, the stemtrac® diagnostic is not reimbursed by the statutory health insurances, but is a self-pay service. Whether and to what extent privately insured patients can receive reimbursement from their insurance company must be clarified with their own private health insurance company.

¹ Pizon, Monika et al. "The number of tumorspheres cultured from peripheral blood is a predictor for presence of metastasis in patients with breast cancer." Oncotarget vol. 7,30 (2016): 48143-48154, doi:10.18632/oncotarget.10174.
² Pizon Monika et al. "Chick Chorioallantoic Membrane (CAM) Assays as a Model of Patient-Derived Xenografts from Circulating Cancer Stem Cells (CCSCs) in Breast Cancer Patients." Cancers vol. 14,6 (2022): 1476. https://doi.org/10.3390/cancers14061476.
³ The stemtrac method is a method produced in the Dr. Pachmann laboratory and is therefore not marketed.